




VLBI Futures, 12 March 2018

# A vision for the 2020s for Reference Frames and Global Astrometry



**Christopher S. Jacobs,** *Jet Propulsion Laboratory, California Institute of Technology*

D. Gordon, A. De Witt, A. Bertarini, C. Garcia-Miro,, S. Horiuchi,

J. McCallum, M. Mercolino, J. Quick, L. Snedeker



Max-Planck-Institut  
für Radioastronomie



HartRAO  
Hartbeespoort Radio  
Astronomy Observatory



European Space Agency




Copyright © 2018 All Rights Reserved. U.S. Government sponsorship acknowledged. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not constitute or imply its endorsement by the United States Government or the Jet Propulsion Laboratory, California Institute of Technology.

# Overview: Current Status of Global Celestial Frames


## VLBI at cm wavelengths:

- SX (8 GHz, 3.6cm) has been only sub-mas frame until last 10 years  
(e.g. *Ma+*, *ICRF1*, 1998, *Ma+*, *ICRF2*, 2009)
- K-band (24 GHz, 1.2cm) now sub-mas (*Lanyi+*, 2010; *de Witt+*, 2016, 2017)
- X/Ka (32 GHz, 9mm) also sub-mas (*Jacobs+*, 2016, 2017)
- Accuracy limited by VLBI systematics due to
  - weak southern geometry: More southern stations?
  - troposphere: WVRs?
  - clocks: distributed clocks?
  - source structure: move to higher frequencies
- Gaia optical: data release #1 is sub-mas for auxiliary quasar solution (*Prusti+*, 2017)  
Final catalog should be sub 100  $\mu$ as for brighter quasars





# SX (8.4 GHz, 3.6cm) *VLBA* + $\sim 100$ other *IVS*




- **Strengths:**
  - 4350 sources
  - Excellent coverage North of  $\delta$  -30 deg
  - median precision  $< 50 \mu\text{as}$
  - SX's 12 million observations, 40 years
  - over 100 stations contributed

- **Weaknesses:**
  - Poor coverage south of  $\delta$  -40 deg
  - only 20% of sources in  $> 10$  sessions
  - source structure worse than K or XKa.




# SX (8.4 GHz, 3.6cm): Dec precision weaker than RA




- **Strengths:**
  - 4350 sources
  - Excellent coverage North of  $\delta$  -30 deg
  - median precision < 50  $\mu$ as
  - SX's 12 million observations, 40 years
  - over 100 stations contributed

- **Weaknesses:**
  - Poor coverage south of  $\delta$  -40 deg
  - only 20% of sources in > 10 sessions
  - source structure worse than K or XKa.





# SX (8.4 GHz, 3.6cm): Dec precision 3X worse RA in south




- **Strengths:**
  - 4350 sources
  - Excellent coverage North of  $\delta$  -30 deg
  - median precision < 50  $\mu$ as
  - SX's 12 million observations, 40 years
  - over 100 stations contributed

- **Weaknesses:**
  - Poor coverage south of  $\delta$  -40 deg
  - only 20% of sources in > 10 sessions
  - source structure worse than K or XKa.




SX: Number Sessions,  $\sim 800 > 10$  sessions, rest 3-5 survey sessions



- **Strengths:**
  - 4350 sources
  - Excellent coverage North of  $\delta$  -30 deg
  - median precision  $< 50 \mu\text{s}$
  - SX's 12 million observations, 40 years
  - over 100 stations contributed

- **Weaknesses:**
  - Poor coverage south of  $\delta$  -40 deg
  - only 20% of sources in  $> 10$  sessions
  - source structure worse than K or XKa.

# K (24 GHz, 1.2cm) *VLBA+ (S. Africa-Tasmania)*




- **Strengths:**
  - Uniform spatial density
  - Galactic plane sources (Petrov+ 2006)
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu\text{as}$
  - needed  $\sim 0.4$  million observations vs. SX's 12 million!

- **Weaknesses:**
  - Ionosphere only partially calibrated by GPS.
  - No solar plasma calibrations
  - South ( $\delta < -30$  deg) weak due to limited HartRAO, South Africa to Hobart, Tasmania data



# K (24 GHz, 1.2cm): Dec precision weaker than RA




- **Strengths:**
  - Uniform spatial density
  - Galactic plane sources (Petrov+ 2006)
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu$ as
  - needed  $\sim 0.4$  million observations vs. SX's 12 million!

- **Weaknesses:**
  - Ionosphere only partially calibrated by GPS.
  - No solar plasma calibrations
  - South ( $\delta < -30$  deg) weak due to limited HartRAO, South Africa to Hobart, Tasmania data



# K (24 GHz, 1.2cm): Dec precision weaker than RA




- **Strengths:**
  - Uniform spatial density
  - Galactic plane sources (Petrov+ 2006)
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu\text{s}$
  - needed  $\sim 0.4$  million observations vs. SX's 12 million!

- **Weaknesses:**
  - Ionosphere only partially calibrated by GPS.
  - No solar plasma calibrations
  - South ( $\delta < -30$  deg) weak due to limited HartRAO, South Africa to Hobart, Tasmania data



## K (24 GHz, 1.2cm): Number sessions 6-9



825 sources

A de Witt et al, 2017

solution D. Gordon 2018 Feb 15


Right Ascension (hours)

Number Sessions

just 1  
just 2  
3 to 5  
6 to 9  
10 to 29  
30 to 69  
70+


- **Strengths:**
  - Uniform spatial density
  - Galactic plane sources (Petrov+ 2006)
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu\text{s}$
  - needed  $\sim 0.4$  million observations vs. SX's 12 million!

- **Weaknesses:**
  - Ionosphere only partially calibrated by GPS.
  - No solar plasma calibrations
  - South ( $\delta < -30$  deg) weak due to limited HartRAO, South Africa to Hobart, Tasmania data



- **Strengths:**
  - Uniform spatial density
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu\text{as}$
  - needed only 70K observations vs. SX's 12 million!


- **Weaknesses:**
  - Poor near Galactic center due to inter-stellar media scattering
  - South weak due to limited time on ESA's Argentina station
  - Limited Argentina-California data makes vulnerable to  $\delta$  zonals
  - Limited Argentina-Australia weakens  $\delta$  from -45 to -60 deg



- **Strengths:**
  - Uniform spatial density
  - less structure than S/X (3.6cm)
  - precision < 100  $\mu$ as
  - needed only 70K observations vs. SX's 12 million!


- **Weaknesses:**
  - Poor near Galactic center due to inter-stellar media scattering
  - South weak due to limited time on ESA's Argentina station
  - Limited Argentina-California data makes vulnerable to  $\delta$  zonals
  - Limited Argentina-Australia weakens  $\delta$  from -45 to -60 deg





- **Strengths:**
  - Uniform spatial density
  - less structure than S/X (3.6cm)
  - precision  $< 100 \mu\text{s}$
  - needed only 70K observations vs. SX's 12 million!

- **Weaknesses:**
  - Poor near Galactic center due to inter-stellar media scattering
  - South weak due to limited time on ESA's Argentina station
  - Limited Argentina-California data makes vulnerable to  $\delta$  zonals
  - Limited Argentina-Australia weakens  $\delta$  from -45 to -60 deg



- **Strengths:**
  - Uniform spatial density
  - less structure than S/X (3.6cm)
  - precision < 100  $\mu$ as
  - needed only 70K observations vs. SX's 12 million!

- **Weaknesses:**
  - Poor near Galactic center due to inter-stellar media scattering
  - South weak due to limited time on ESA's Argentina station
  - Limited Argentina-California data makes vulnerable to  $\delta$  zonals
  - Limited Argentina-Australia weakens  $\delta$  from -45 to -60 deg



# Global Astrometry: path to progress

- **Source structure:**

Source more compact at higher frequency

Move astrometry to K or XKa

- **Troposphere:** Wet troposphere fluctuations are major error

Build in WVRs

- **Clocks: Achieve common clocks**

distribute clocks to southwestern stations

then continental stations. Technical Feasibility? Method: fiber? Cost?

- **Geometry/Resolution:**

VLBA does very well in the northern hemisphere

but at far end of reach  $\delta \sim -30$  deg, loses accuracy.

Astrometry would benefit from longer N-S baselines:

southern outriggers?

VLBA antennas in Mexico?


at ALMA site?

- **Sensitivity:**

Current systems use 0.5 GHz analog @ 2bit = 2 Gbps

Large antennas are expensive, so gain sensitivity via increased bandwidth: 8 to 32 Gbps?



# Radio Source Structure vs. Frequency



**X-band**  
**8.6 GHz**  
**3.6cm**

**K-band**  
**24 GHz**  
**1.2cm**

**Q-band**  
**43 GHz**  
**0.7cm**



**The sources**  
**become better →**  
**Less structure**

**Ka-band**  
**32 GHz**  
**0.9cm**

Images credit: Pushkarev & Kovalev *A&A*, 544, 2012 (SX);

Charlot et al, *AJ*, 139, 2010 (KQ)



# Calibrating Troposphere Turbulence

- JPL Advanced Water Vapor Radiometer

- ~ 1 deg beam better matches VLBI
  - improved gain stability
  - improved conversion of brightness temperature to path delay

Tanner & Riley, Radio Sci., 38, 2003

<http://adsabs.harvard.edu/abs/2003RaSc...38.8050T>



- Initial demos show 1mm accuracy  
Goldstone-Madrid 8000 km baseline  
using X/Ka phase delays


*Jacobs et al, AAS Winter 2005.*

*Bar Sever et al, IEEE, 2007.*

<http://adsabs.harvard.edu/abs/2007IEEEP..95.2180B>

- A-WVRs deployed at Goldstone/Madrid  
Seeking funding for Tidbinbilla, Aus
- A-WVR not used yet for Operations

## VLBI Delay Residuals DOY 200 Ka-Band DSS26-DSS55





# NASA/ESA Deep Space Net Geometry




ESA Argentina to NASA-California under-observed by order of magnitude!

## Baseline percentages

- Argentina is part of 3/5 baselines or 60%  
but only 13% of obs
- Aust- Argentina 7.5%
- Spain-Argentina 2.2%
- Calif- Argentina 3.7%

This baseline is under-observed by a factor of ~ 12.

More time on ESA's Argentina station would have a huge, immediate impact!!



Maps credit: Google maps


- ESA's Argentina 35-meter antenna **adds 3 baselines** to DSN's 2 baselines
- Full sky coverage by accessing south polar cap
  - near perpendicular mid-latitude baselines: CA to Aust./Argentina

### Zonal Errors

- $\Delta\text{RA}$  vs. Dec:  
~300  $\mu\text{s}$  in south, 200  $\mu\text{s}$  in north
- Need 2 baselines to get 2 angles:  
California-Canberra: 24K obs  
California-Argentina: 2K obs
- > Need more California-Argentina data to overcome this 12 to 1 distortion in sampling geometry.  
ESA's Malargüe is key.
- Usuda, Japan 54-m XKa (2019) would improve North-South sampling geometry and thus control declination zonal differences.



### XKa vs. SX: Zonal errors





# Sensitivity

- **Integrations are typically 1-2 minutes**

Driven by

Need to monitor large number of sources

Coherence time is short at 24 and 32 GHz

- **Data rates**

VLBA-only 2 Gbps SX and K

DSN 2 Gbps (0.256 for Argentina)

IVS 0.256 to 0.512 Gbps for large networks

Limits much reference frame work to Fluxes  $> 100$  mJy

- **Future:**


Increase data rates to 8 to 32 Gbps or more?

Allows access to weaker sources and/or faster scans




# Optical vs. Radio Frames

Systematics to be flushed out via  
Inter-comparison of multiple high  
precision frames.



Credit: Marscher+, Krichbaum+



## Systematics:

Gaia: 60 mas beam sees  
Host galaxy, foreground stars, etc.

ALMA: pilot obs bright end  $\sim 5^{\text{mag}}$   
Waiting on 10km+ configurations

VLBI: All bands need more  
southern data

S/X: Source structure

K: Ionosphere

XKa: Argentina baselines  
under-observed

# Tying optical and Radio Celestial Frames

## Gaia DR1-aux vs. VLBI



|   | SX-band<br>8 GHz<br>3.6cm | K-band<br>24 GHz<br>1.2 cm | XKa-band<br>32 GHz<br>0.9 cm |
|---|---------------------------|----------------------------|------------------------------|
| # Observations                                      | 12 million                | 0.25 million               | 0.06 million                 |
| # sources   | 1926                      | 473                        | 405                          |
| # outliers $> 5\sigma$                              | 100                       | 13                         | 6                            |
| % outliers  | 5.2 %                     | 2.7 %                      | 1.5 %                        |
| $\alpha$ wRMS                                       | 523 $\mu$ as              | 431 $\mu$ as               | 433 $\mu$ as                 |
| $\delta$ wRMS                                       | 531 $\mu$ as              | 453 $\mu$ as               | 418 $\mu$ as                 |
| $R_x$   | -37 +- 13                 | -89 +- 24                  | 57 +- 24                     |
| $R_y$   | 0 +- 11                   | 14 +- 21                   | 32 +- 21                     |
| $R_z$   | -29 +- 13                 | -13 +- 23                  | 21 +- 24                     |
| $\Delta\alpha$ vs. $\delta$ tilt<br>( $\mu$ as/deg) | -0.46 +- 0.25             | -1.55 +- 0.53              | -2.83 +- 0.58                |

Rx vulnerable  
To trop errors

Hints that results improve by going to higher radio frequency

However, the above results do not use exact same objects (Jacobs+, IAU 330, 2017)




# Summary: Global Astrometry

- **Current precisions  $\sim 100 \mu\text{as}$  or better at three bands:**
  - SX 8 GHz 4200 objects, **S-band RFI source structure**
  - K 24 GHz 825 objects, **no dual band in cals (C/K?)**
  - XKa 32 GHz 680 objects, **not supported by VLBA**
- **Source structure:** Move astrometry to K or Xka
- **Troposphere:** Built in WVRs
- **Clocks:** Phased plan to distribute clocks to southwestern stations then continental stations
- **Geometry/Resolution:**
  - VLBA does very well in the northern hemisphere but at far end of reach  $\delta \sim -30^\circ$  deg, loses accuracy.
  - Astrometry would benefit from longer N-S baselines: southern outriggers?**
  - VLBA antennas in Mexico? at ALMA site?**
- **Sensitivity:**
  - Current systems use 0.5 GHz analog @ 2bit = 2 Gbps
  - Large antennas are expensive, so gain sensitivity via increased bandwidth: 8 -32 Gbps?

# Backup



# Active Galactic Nuclei (*Marscher*)



$R \sim 0.1 - 1 \mu\text{as}$

1 mas

Features of AGN: *Note the Logarithmic length scale.*



“Shock waves are frequency stratified, with highest synchrotron frequencies emitted only close to the shock front where electrons are energized. The part of the jet interior to the mm-wave core is opaque at cm wavelengths. At this point, it is not clear whether substantial emission occurs between the base of the jet and the mm-wave core.”

*Credits: Alan Marscher, 'Relativistic Jets in Active Galactic Nuclei and their relationship to the Central Engine,' Proc. of Science, VI Microquasar Workshop: Microquasars & Beyond, Societa del Casino, Como, Italy, 18-22 Sep 2006. Overlay (not to scale): 3 mm radio image of the blazar 3C454.3 (Krichbaum et al. 1999)*



# SX VLBI systematic Floor $\sim 20$ to $30 \mu\text{as}$ ?



## Set of Flicker Noise sources



## Set of White Noise sources



Green ★ : ICRF2 noise floor - average on sources in  $15^\circ$  declination bands.



Attention! This method uses ALL “good” sessions, contrary to the decimation test.

Le Bail+ (EVGA, 2017) use Allan variance test on position time histories to determine when **averaging no longer helps—systematic floor is encountered.**

**Structure part of this floor should be several times smaller at K (24 GHz) and Ka (32 GHz)**

# Optical vs. Radio systematics offsets



## SDSS Optical images of quasars (scale 5-10 asec)



Credit: SDSS

- Optical structure: The host galaxy may not be centered on the AGN or may be asymmetric.
- Optical systematics unknown, fraction of milliarcsecond optical centroid offset?
- Optical imaging generally 10s of milliarcsecond. In general, no sub-mas optical imaging.

# Optical vs. Radio systematics offsets



Petrov & Kovalev (MNRAS, 2017) show that optical-radio astrometric offsets Correlate with jet direction (or anti-direction).

They argue that the offsets are dominated by optical synchrotron jets.




# Optical vs. Radio systematics offsets



Petrov & Kovalev (MNRAS, 2017)

- Example of optical jet in “nearby” 3C 264 would scale to ~milli-arcsecond offsets at typical AGN distances.
- Optical synchrotron jets may be limiting factor in radio-optical astrometric agreement.
- VLBI interferometry “locks” onto the brightest component. Also extremely high resolution resolves out extended structures. So VLBI positions is close to the core.
- Gaia optical image’s centroid averages all of the light distribution, jet included. “Beam” is 60 milliarcseconds.
- Optical may be more easily biased than radio.



**Figure 3.** The archival HST image of 3C264 at 606 nm, HST project ID 13327 (Meyer et al. 2015).